



WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

PCT

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification<sup>7</sup> :

C12N 9/00

A2

(11) International Publication Number:

WO 00/06706

(43) International Publication Date: 10 February 2000 (10.02.00)

(21) International Application Number:

PCT/EP99/05413

(22) International Filing Date:

28 July 1999 (28.07.99)

(30) Priority Data:  
9816639.0

30 July 1998 (30.07.98)

GB

(71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).

(71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESellschaft MBH [AT/AT]; Brunnerstrasse 59, A-1230 Vienna (AT).

(72) Inventors; and  
(75) Inventors/Applicants (for US only): REVENKOVA, Ekaterina [RU/CH]; Peter Rot-Strasse 76, CH-4058 Basel (CH); PASZOWSKI, Jerzy [PL/CH]; Blauenweg 10, CH-4224 Menzingen (CH).

(74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).

(54) Title: MAP KINASE PHOSPHATASE MUTANT

(57) Abstract

The present invention relates to DNA encoding proteins contributing to the regulation of a plant's response to DNA damage. DNA according to the invention comprises an open reading frame encoding a protein characterized by a stretch of amino acids or component amino acid sequence having 40% or more identity with an aligned component sequence of SEQ ID NO:3. Preferably the DNA encodes a MAP kinase phosphatase.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MR	Mauritania	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	VU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China			PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

MAP kinase phosphatase mutant

The present invention relates to DNA encoding proteins contributing to the regulation of a plant's response to abiotic stress and in particular genotoxic stress.

Cells of all organisms have evolved a series of DNA repair pathways which counteract the deleterious effects of DNA damage and are triggered by intricate signal cascades. To be able to modify or improve DNA repair using gene technology it is necessary to identify key proteins involved in said pathways or cascades. Therefore it is the main object of the present invention to provide DNA comprising an open reading frame encoding such a key protein.

DNA according to the present invention comprises an open reading frame encoding a protein characterized by a stretch of amino acids or component amino acid sequence having 40% or more identity with an aligned component sequence of SEQ ID NO: 3. The protein characterized by SEQ ID NO: 3 is tracked down with the help of a T-DNA tagged Arabidopsis mutant showing hypersensitivity to methyl methanesulfonate (MMS). Said hypersensitivity as well as an observed hypersensitivity to other DNA damaging treatments such as UV light is indicative of the proteins' involvement in the repair of DNA damage, or in signaling pathways implicated in the response to similar genotoxic stress. The mutant is also sensitive to elevated temperature and anti-oxidant N-acetylcysteine. The mutant is not sensitive to osmotic shock, increased salinity, oxidative stress or elevated ethylene levels. An important characteristic of the mutant is cell death in response to growth in small closed vessels. This phenotype can be complemented by addition of abscisic acid (ABA) to the growth media. Furthermore, the mutant is more sensitive to exogenously applied ABA compared with the wild type which supports the notion that the genes disclosed by the present invention (SEQ ID NO: 1) are involved in stress signaling mediated by ABA.

Sequence alignments of SEQ ID NO: 3 using commercially available computer programs such as BLASTP of the NCBI BLAST family of programs or TFASTA or BestFit of the Wisconsin Package Software, all based on well known algorithms for sequence identity or similarity searches, reveal that stretches of SEQ ID NO: 3 (component sequences) having more than 100 and preferably between 120 to 250 amino acids length can show between

20% and almost 40% sequence identity to aligned stretches of known phosphatases, particularly phosphotyrosine phosphatases, MAP kinase phosphatases or dual specificity phosphatases. Protein phosphatases are classified by their substrate specificities as either phosphoserine/threonine phosphatases (PSTPs) or phosphotyrosine phosphatases (PTPs). The dual specificity phosphatases (DSPs) dephosphorylate both phosphotyrosine and phosphoserine/threonine residues and represent a subfamily of PTPs. MAP kinase phosphatases (MKPs) belong to the family of DSPs. The sequence VHCCQGVSR (SEQ ID NO: 4) found in SEQ ID NO: 3 can be interpreted as corresponding to the mammalian sequence motif IHGXAGXXRS (SEQ ID NO: 5) defining the family of PTPs, wherein the Ile at the first position can be replaced by Val and the Ser at the last position can be replaced by Thr

The present invention defines a new protein family the members of which are characterized by component amino acid sequences of more than 100 amino acid length showing 40% or higher amino acid sequence identity to aligned component sequences of SEQ ID NO: 3. Preferably said component sequences are of more than 120, more than 160 or even more than 200 amino acids length. The amino acid sequence identity is preferably higher than 50% or even higher than 55%. Most preferred are identities higher than 70%.

An example of DNA according to the present invention is described in SEQ ID NO: 1. The amino acid sequence of the protein encoded is identical to SEQ ID NO: 3. After alignment a stretch of the protein having about 140 amino acids shows 36% sequence identity to the MKP-1 protein described by Sun et al (Cell 75: 487-493, 1993). The identity determined after alignment with MKP-2 and MKP-3 is determined as 34% and 26%, respectively. Thus, according to the present invention a protein family related to MAP kinase phosphatases can be defined the members of which after alignment of a stretch of more than 100 amino acids length show 40% or higher amino acid sequence identity to SEQ ID NO: 3. Preferably, the amino acid sequence identity is higher than 50% or even higher than 55%. When making multiple sequence alignments, certain algorithms can take into account sequence similarities, such as same net charge or comparable hydrophobicity/hydrophilicity of the individual amino acids, in addition to sequence identities. The resulting values of sequence similarities, as compared to sequence identities, can help to assign a protein to the correct protein family in border-line cases. Proteins of particular interest, within the scope of the present invention, are MAP kinase phosphatases the amino acid sequence of which comprises at least one of the following characteristic amino acid subsequences:

- (a) TSILYDVFDYFEDV (SEQ ID NO: 6)
- (b) FVHCQGVSRST (SEQ ID NO: 7)
- (c) FVHC (SEQ ID NO: 8)
- (d) QGVSR (SEQ ID NO: 9)
- (e) YFKSD (SEQ ID NO: 10)

DNA encoding proteins belonging to the new protein family according to the present invention can be isolated from monocotyledonous and dicotyledonous plants. Preferred sources are corn, sugar beet, sunflower, winter oilseed rape, soybean, cotton, wheat, rice, potato, broccoli, cauliflower, cabbage, cucumber, sweet corn, daikon, garden beans, lettuce, melon, pepper, squash, tomato, or watermelon. However, they can also be isolated from mammalian sources such as mouse or human tissues. The following general method, can be used, which the person skilled in the art will normally adapt to his specific task. A single stranded fragment of SEQ ID NO: 1 or SEQ ID NO: 2 consisting of at least 15, preferably 20 to 30 or even more than 100 consecutive nucleotides is used as a probe to screen a DNA library for clones hybridizing to said fragment. The factors to be observed for hybridization are described in Sambrook et al, Molecular cloning: A laboratory manual, Cold Spring Harbor Laboratory Press, chapters 9.47-9.57 and 11.45-11.49, 1989. Hybridizing clones are sequenced and DNA of clones comprising a complete coding region encoding a protein with more than 40% sequence identity to SEQ ID NO: 3 is purified. Said DNA can then be further processed by a number of routine recombinant DNA techniques such as restriction enzyme digestion, ligation, or polymerase chain reaction analysis.

The disclosure of SEQ ID NO: 1 enables a person skilled in the art to design oligonucleotides for polymerase chain reactions which attempt to amplify DNA fragments from templates comprising a sequence of nucleotides characterized by any continuous sequence of 15 and preferably 20 to 30 or more basepairs in SEQ ID NO: 1. Said nucleotides comprise a sequence of nucleotides which represents 15 and preferably 20 to 30 or more basepairs of SEQ ID NO: 1. Polymerase chain reactions performed using at least one such oligonucleotide and their amplification products constitute another embodiment of the present invention.

Knowing the nucleotide sequence of the *Arabidopsis* MKP1 gene and the amino acid sequence of the encoded protein it is possible to identify proteins interacting with AtMKP1 and to clone their corresponding genes using well known techniques. For example radioactively labeled AtMKP1 protein can be used for interactive cloning on cDNA expression libraries. AtMKP1 protein or parts thereof can be used to generate polyclonal or monoclonal antibodies specific for AtMKP1. The *AtMKP1* gene can be used to generate variants of AtMKP1 protein tagged with GST, MYK or His. Said antibodies and MKP1 variants allow to isolate native protein complexes by immunoprecipitation and to determine sequences of proteins present in these complexes by micro-sequencing. The resulting sequence information can in turn be used to clone corresponding genes. Alternatively, said antibodies or tagged MKP1 variants can be used to screen epitope libraries for epitopes which interact with AtMKP1 protein. The AtMKP1 protein and parts thereof, in particular the N-terminal 490 amino acid region and the C-terminal 492 amino acid region can also be used to search for interacting proteins with a Two-hybrid system (e.g. in yeast, in mammalian cells, or in bacteria). This allows to obtain sequence information about interacting proteins.

Based on the disclosed finding that AtMKP1 proteins are involved in a plant's abiotic environmental stress response, it becomes possible to engineer the corresponding signaling pathway, of which AtMKP1 is a part, to be chemically regulated due to chemical activation or repression of transgenes encoding AtMKP1 or proteins interacting therewith. Such plants can be obtained by transformation with the corresponding genes under control of chemically inducible promoters. Application of inducers is expected to modify the activity of the AtMKP1 signaling pathway and to result in altered adaptation to abiotic environmental stress. Alternatively, AtMKP1 protein or its interacting proteins can be used as targets for chemicals inhibiting or stimulating their activities which again is expected to modify abiotic stress responses.

## EXAMPLES:

### Example 1: *Cloning of the gene responsible for the mkp1 mutant phenotype*

*Arabidopsis* T-DNA insertion lines as produced by the INRA-Versailles and available from the Nottingham Arabidopsis Stock Center (NASC) are screened for sensitivity to methylmethanesulfonate (MMS) at a concentration of 100 ppm as described by Masson et al.

Genetics 146: 401-407, 1997. Plants which die in the presence of 100 ppm MMS are found in the family AAN4. Thus, the corresponding T-DNA insertion mutation is assumed to give rise to this hypersensitive phenotype. This assumption is supported by genetic analysis showing co-segregation of the hypersensitive phenotype with the T-DNA insertion.

Genomic DNA from the mutant plants is isolated as described by Dellaporta et al, Plant Mol Biol Reporter 1: 19-21, 1983. A fragment of genomic DNA flanking the right border of the inserted T-DNA is rescued essentially according to Bouchez et al, Plant Mol Biol Reporter 14: 115-123, 1996, with minor modifications. Genomic DNA is digested with PstI, ethanol precipitated and resuspended in H<sub>2</sub>O. DNA of vector pResc38 (Bouchez et al supra) is digested with PstI and dephosphorylated with shrimp alkaline phosphatase. The phosphatase is heat inactivated, the vector DNA is ethanol precipitated and resuspended in H<sub>2</sub>O. 2.5 µg of PstI digested genomic DNA and 2.5 µg of PstI digested and dephosphorylated vector are mixed and ligated overnight at room temperature in a total volume 100 µl in the presence of 10 units of T4 DNA ligase. The DNA of the ligation mixture is precipitated with ethanol, resuspended in 50 µl H<sub>2</sub>O, and digested with XbaI in a total volume of 100 µl. XbaI digested DNA is precipitated with ethanol and resuspended in H<sub>2</sub>O. A second overnight ligation reaction in the presence of 10 units T4 DNA ligase is performed in a total volume of 200 µl at room temperature to achieve circularization of DNA fragments. The DNA of the ligation mixture is again precipitated with ethanol, rinsed two times with 70% ethanol, dried and dissolved in 5 µl H<sub>2</sub>O. Two 2 µl aliquots are used for electroporation of electrocompetent E.coli XL1-Blue cells (Stratagene) according to the manufacturer's instructions. Transformants containing the inserted T-DNA and adjacent *Arabidopsis* genomic DNA sequences are selected on plates with 50mg/l kanamycin. Single bacterial colonies are analyzed by isolation of plasmid DNA using QIAprep Spin Plasmid Kit (Qiagen) and restriction digestion with PstI and XbaI. Plasmid pBN1 containing 3.7 kb of inserted T-DNA linked to 5 kb of *Arabidopsis* DNA is identified. Sequencing of the junction site is performed using a primer directed towards the flanking plant DNA and having the nucleotide sequence 5' -GGTTCTACAGGACGTAACAT-3' (SEQ ID NO: 14) complementary to T-DNA 41 nucleotides from the right border. Digestion of this clone with SstI allows isolation of a 960 bp fragment which when labelled with <sup>32</sup>P can be conveniently used as a probe to screen wild type *Arabidopsis* genomic and cDNA libraries in order to identify the wild type gene affected in the *mkp1* mutant line.

**Example 2: Cloning of the AtMKP1 wild-type gene**

The 960 bp SstI fragment mentioned at the end of example 1 is labeled with  $^{32}\text{P}$  by random oligonucleotide-primed synthesis (Feinberg et al, Anal Biochem 132: 6-13, 1983) for use as a probe in the following hybridization experiments.

Southern blot analysis of *Arabidopsis* wild type and *mkp1* DNA digested with EcoRV confirms that in the *mkp1* genomic DNA the sequence hybridizing to the probe is linked to T-DNA.

Northern blot analysis of *Arabidopsis* wild type RNA reveals the presence of a hybridizing transcript in RNA extracted from seven-day-old wild type seedlings. No such hybridizing fragment is detected in the corresponding RNA of *mkp1* seedlings.

A cDNA library (Ellledge et al, 1991) and a genomic library (Stratagene) of wild type *Arabidopsis thaliana* ecotype Columbia is screened with the labelled SstI fragment

mentioned above. Screening of the bacteriophage  $\lambda$  libraries is performed according to the protocols described in chapter 6 of Ausubel et al, 1994, "Current protocols in molecular biology", John Wiley & Sons, Inc. Hybridization is performed as described by Church and Gilbert, Proc Natl Acad Sci USA 81: 1991-1995, 1984. Bacteriophage clones hybridizing to SstI fragment are subjected to *in vivo* excision of plasmids according to Ellledge et al, Proc Natl Acad Sci USA 88: 1731-1735, 1991, and Stratagene protocols. Inserts of the plasmids obtained are further analyzed by sequencing.

By partial sequencing and alignment of ten overlapping clones (pBN5.1 to pBN5.10) isolated from the genomic library a continuous genomic sequence of 6356 bp (see SEQ ID NO: 1) is decoded.

Ten cDNA clones representing the same gene, one of them a 3.0 kb full-length cDNA (SEQ ID NO: 2), are isolated from the cDNA library.

**Example 3: Sequence Analysis and Alignments**

The 3 kb full-length cDNA clone of SEQ ID NO: 2 encodes an ORF with the start codon being defined by basepairs 298-300 and the stop codon by basepairs 2650-2652. The ORF encodes a protein consisting of 784 amino acids (SEQ ID NO: 3) and a predicted molecular mass of 86.0 kD. Alignment with the genomic sequence of SEQ ID NO: 1 reveals three introns. T-DNA is inserted within the coding sequence of the *mkp1* mutant DNA before basepair position 502 according to the numbering of SEQ ID NO: 2. The sub-sequence



VHCCGGVSRS (SEQ ID NO: 4) found in SEQ ID NO: 3 can be interpreted as

corresponding to the mammalian sequence motif IHGXAGXXRS (SEQ ID NO: 5) defining

the family of protein tyrosine phosphatases, wherein the Ile at the first position can be

replaced by Val and the Ser at the last position can be replaced by Thr (Van Vactor et al,

Curr Opin Gen Dev 8: 112-126, 1998). Therefore it is concluded that the wild type ORF

encodes a protein tyrosine phosphatase with invariant aspartic acid, cysteine, and arginine

residues (Fauman et al, Trends Biochem Sci 21: 413-417, 1996) in positions 204, 235 and

241 according to SEQ ID NO: 3.

A data base search using the TFASTA program (Wisconsin Package Version 9.1, Genetics

Computer Group (GCG), Madison, Wisc.) reveals that the encoded phosphatase has a

significant similarity to dual specificity phosphatases. The closest homologue identified is

*Xenopus laevis* MAP kinase phosphatase (MKP; Lewis et al, J Cell Sci 108: 2885-2896,

1995) showing 38.1% identity and 52.5% similarity in a 140 amino acid overlap region. The

deduced AtMKP1 protein also has 36.0% identity and 52.5% similarity with a 140 amino

acid overlap region encoded by the rat 3CH134/CL100 cDNA representing a rat MKP1.

Essentially identical results are obtained when using the BLASTP 2.0.4 (Feb-24-1998)

program of the NCBI BLAST family of programs which, allowing gapped alignment,

compares an amino acid query sequence against a protein sequence database (Altschul et

al, Nucleic Acids Res. 25: 3389-3402, 1997). No higher plant homologues are identified.

The genomic position of the AtMKP1 gene is determined by hybridization to filters containing genomic YAC clones publicly available from the Arabidopsis Biological Resource Center (Ohio, USA). The AtMKP1 gene is found to map to chromosome 3 between markers ve022 and BGL1.

#### Example 4: *Complementation*

mkp1 mutant plants are transformed with DNA comprising the corresponding wild type

genomic DNA including promoter and polyadenylation signal to find out whether the cloned

wild type gene is able to complement the mutant mkp1 phenotype.

mkp1 mutant plants harbor T-DNA containing the *NPTII* and *bar* marker genes under the

control of nos and CaMV35S promoters, respectively. Therefore, different marker genes are

used for the transformation construct. The vector used is a derivative of p1'barbi which is

highly efficient in *Arabidopsis* transformation (Mengiste et al, Plant J 12: 945-948, 1997). In

p1'barbi the EcoRI fragment containing 1'promoter, bar gene coding region, and CaMV 35S polyadenylation signal is inverted in respect to the T-DNA borders by EcoRI digestion and re-ligation. In the resulting plasmid the 1'promoter (Velten et al, EMBO J 3: 2723-2730, 1984) is directed towards the right border of the T-DNA. This plasmid is digested with BamHI and NheI, and the bar gene and CaMV 35S polyadenylation signal are replaced by a synthetic polylinker with the sites for the restriction enzymes BamHI, HpaI, ClaI, StuI and NheI. The resulting plasmid is digested with BamHI and HpaI and ligated to a BamHI-PvuII fragment of pROB1 (Bilang et al, Gene 100: 247-250, 1991) containing the hygromycin-B-resistance gene *hph* linked to the CaMV 35S polyadenylation signal. The T-DNA of the resulting binary vector p1'hygi contains the hygromycin resistance selectable marker gene under the control of the 1'promoter and unique cloning sites for the restriction enzymes *ClaI*, *StuI* and *NheI* located between the marker gene and the T-DNA right border. p1'hygi is used to insert the reconstructed *AtMKP1* gene as follows. Plasmid pBN1 of example 1 is digested with *PstI* and *MunI* and dephosphorylated. The restriction fragment containing the 3'portion of the *AtMKP1* gene and pBluescript-SK(+) is purified from the agarose gel and ligated to the *PstI*-*MunI* restriction fragment of the wild type genomic clone pBN5.2 (example 2) including the 5' end of the coding sequence of the *AtMKP1* gene and 2.4 kb of upstream sequences. The reconstructed *AtMKP1* gene is excised by *PstI* and *NotI* and after filling the ends is inserted into the *StuI* site of p1'hygi. The construct is introduced by transformation into *Agrobacterium tumefaciens* strain C58CIRif<sup>R</sup> containing the non-oncogenic Ti plasmid pGV3101 (Van Larebeke et al, Nature 252: 169-170, 1974). T-DNA containing the reconstructed *AtMKP1* gene is transferred to mutant plants by the method of *in planta Agrobacterium* mediated gene transfer (Bechthold et al, C R Acad Sci Paris, Life Sci 316: 1194-1199, 1993). Seeds of infiltrated plants are grown on hygromycin-containing medium to screen for transformants. The progeny of selfed hygromycin resistant plants is analyzed for the segregation of hygromycin resistance. The families in which a 3:1 segregation ratio is observed are used to isolate homozygous lines bearing the newly introduced T-DNA inserted at a single genetic locus. The obtained hygromycin resistant lines are analyzed by Northern blot analysis for the restoration of *AtMKP1* expression. In these lines the restoration of transcription of the *AtMKP1* gene can be observed as well as the restoration of the wild type level of MMS resistance and ABA mediated stress responses. Complementation is not observed in plants transformed with p1'hygi only.

### Example 5: Cloning of homologous sequences from other plant species

Use of *AtMKP1* cDNA as a probe for Southern hybridization with genomic DNA from other plant species such as *Sinapis alba* (mustard), *Lycopersicum esculentum* (tomato) and *Zea mays* (maize) is successful in the case of *Sinapis alba* which belongs to the same family as *Arabidopsis* (Brassicaceae).

Homologous sequences from the other species can be identified in a PCR approach using degenerate primers 1-3 below, wherein I is inosine, derived from the regions conserved between VH-PTP13 of *Clamydomonas eugametos* and *AtMKP1* protein:

Primer 1 (forward): 5' - AAY AAY GGI ATH ACI CAY ATH YT-3' (SEQ ID NO: 11);

Primer 2 (reverse): 5' - YTG RCA IGC RAA ICC CAT RTT IGG-3' (SEQ ID NO: 12);

Primer 3 (reverse): 5' - IGT CCA CAT IAR RTA IGC DAT IAC (SEQ ID NO: 13);

A PCR reaction is performed in a total volume of 50 µl containing 1x reaction buffer (Qiagen), 200 µM of each dNTP, 1.25 units of Taq polymerase (Qiagen), and 100 pmol of each primer.

Reaction 1 is performed with primers 1 and 2 using genomic DNA from *Sinapis alba* (200 ng), *Lycopersicum esculentum* (400 ng), or *Zea mays* (600 ng) as the original template DNA. Amplification is carried out after an initial denaturation step of 3 min at 94°C, followed by 30 cycles of 30 sec at 94°C, 30 sec at 40°C, and 3 min at 72°C. The resulting amplification mixture is diluted 10<sup>3</sup> fold.

Reaction 2 is performed using 2 µl of the above dilution to provide the necessary template DNA. This time primers 1 and 3 are used under the same conditions as specified for reaction 1. The resulting amplification products are cloned into the T/A vector pCR2.1 (Invitrogen) and further analyzed by nucleotide sequencing.

Using this PCR approach it is possible to amplify sequences homologous to the *AtMKP1* gene from all the species mentioned above. Whereas the nucleotide sequence from *Sinapis alba* *SamKP1* (SEQ ID NO: 15 encoding SEQ ID NO: 16) is 90.8% identical to the *AtMKP1* sequence, the nucleotide sequence from *Lycopersicum esculentum* *LeMKP1* (SEQ ID NO: 17 encoding SEQ ID NO: 18) is 72.3% and the *Zea mays* sequence

*ZmMKP1* (SEQ ID NO: 19 encoding SEQ ID NO: 20) 71.8% identical. The fragments hybridize to genomic DNA from corresponding species under the usual hybridization conditions for Southern blot analysis. The fragments can be used as probes to screen cDNA libraries for corresponding cDNA sequences.

The 243 bp *ZmMKP1* fragment amplifying from maize DNA is used as a probe to screen a maize cDNA library (Clontech) made in the Lambda ZAP<sup>®</sup> II Vector (Clontech) from "Blizzard" hybrid etiolated shoots, which were treated with the herbicide safener Benoxacor. The titer of the library is determined as  $3 \times 10^9$  pfu/ml.

Library screening is conducted as described in the Clontech Lambda Library Protocol Handbook, with some slight modifications. Briefly, a single colony of XL-1 Blue is picked and incubated overnight at 37°C in LB medium, containing 10mM MgSO<sub>4</sub> and 0.2% maltose. 600 µl of stationary phase grown bacteria for each 150 mm plate is combined with 100 µl of phage library dilution in sterile 1× lambda dilution buffer (100mM NaCl; 10mM MgSO<sub>4</sub>; 35mM Tris-HCl, pH7.5) to yield approximately 30,000 pfu per plate. This mixture is incubated at 37°C for 15 minutes, subsequently 7 ml of melted LB soft top agarose (at 48°C) is added to the cell suspension for each 150 mm plate, shortly mixed and then poured on two-day-old LB<sup>MgSO<sub>4</sub></sup> agar plates, which have been pre-warmed to 37°C for four hours. The plates are then incubated at 37°C until plaques reach appropriate sizes (after about 8 to 9 hours). After chilling the plates at 4°C for one hour, phage particles are transferred to Hybond N nitrocellulose filters and the orientation of each filter to its plate is recorded with a waterproof pen. The filters are then treated by placing them on Whatman 3MM paper saturated with the appropriate solution. The treatments include denaturation solution (0.5M NaOH; 1.5M NaCl) for 2 minutes, followed by neutralization solution (0.5M Tris-HCl, pH 7.2; 1.5M NaCl; 1mM EDTA) for 3 minutes and 2× SSC for 3 minutes. DNA is subsequently crosslinked to the filters by UV.

Filters are then pre-hybridized, hybridized with the radioactively labeled *ZmMKP1* fragment and washed as described in Sambrook et al, Molecular cloning: A laboratory manual, Cold Spring Harbor Laboratory Press, chapters 9.47-9.57 and 11.45-11.49, 1989. An agar plug from the position of a positive plaque is then removed from the master plate and incubated overnight at 4°C in 1 ml of 1× lambda dilution buffer, containing 20 µl of chloroform. Each filter is determined and the phages are re-plated to obtain approximately 200 to 500 plaques on a 150 mm plate for a secondary screen as described above. Single plaques of interest

are collected from the agar plates and incubated over night at 4°C in 500 µl of 1× lambda dilution buffer and 20 µl of chloroform.

The Bluescript phagemid is excised from the λZAP™ vector as described by the *In Vivo* Excision Protocol using the ExAssist/SOLR System in the Stratagene Uni-ZAP™ XR Library Instruction Manual (1993). A 1/100 dilution is made of XL1-Blue MRF' and SOLR overnight cultures (at 30°C) and incubated at 37°C for 2-3 hours. XL1-Blue MRF' cells are then pelleted for 10 minutes at 1,500×g and re-suspended at an OD<sub>600</sub> = 1.0 in 10mM MgSO<sub>4</sub>. 200 µl of these XL1-Blue cells are then combined with 250 µl of phage stock and 1 µl of ExAssist helper phage in a 50 ml conical tube and incubated at 37°C for 15 minutes. 3 ml of LB broth is added and incubated at 37°C for 5 hours, after which the cells are pelleted for 15 minutes at 2,000×g and the supernatant transferred to a new tube. The tube is then heated at 70°C for 15 minutes and centrifuged again for 15 minutes at 4,000×g. The supernatant, containing the excised phagemid pBluescript packaged as filamentous phage particles, is decanted into a new tube. 10 and 100 µl of this phage stock are then added to two tubes with 200 µl of SOLR cells that have been allowed to grow to OD<sub>600</sub> = 0.5-1.0 before being removed from the incubator and further incubated at room temperature. The tubes are incubated at 37°C for 15 minutes, followed by plating 10-50 µl from each tube on LB<sup>amp</sup> (50 µg/ml) and over night incubation at 37°C.

The positive clones are checked for insert size by *EcoRI*/*XhoI* double digestion and end-sequencing with T3 and T7 promoter primers (Promega).

Screening of 360,000 pfu of the library results in three identical clones of 2.2 kb containing the 3' poly(A) tail, but lacking part of the 5' end, including the translation initiation site. The gene corresponding to the identified partial cDNA clone is named *ZmMKP2*, as it is not identical with the *ZmMKP1* fragment used as the probe (92.3% identity on the nucleotide level over the 196 bp fragment flanked by the primers 1 and 3). An additional 213 nucleotides are amplified and cloned by 5' RACE (rapid amplification of cDNA ends) carried out following the instructions of the 5'/3' RACE Kit (Boehringer Mannheim) resulting in a longer cDNA sequence of 2,452 bp but still not complete, judged by the predicted mRNA length from the RNA gel blot analysis and the absence of a possible translation initiation site.

The sequence information gained from the *ZmMKP2* cDNA including the additional 213 nucleotides obtained by 5' RACE (SEQ ID NO: 21 encoding SEQ ID NO: 22) is used to

design two additional backward oriented degenerate primers wherein 1 is inosine to 3' regions conserved between the deduced peptide sequences of *ZmMKP2* and *AtMKP1*:

Primer 4 (reverse): 5'-GCI GCY TTI GCR TCY TTY TCC-3' (SEQ ID NO: 25);

Primer 5 (reverse): 5'-YTC ICK IGC IGG IAR RTG IGT YTC-3' (SEQ ID NO: 26)

These primers are used to PCR amplify a larger fragment of a MAP kinase phosphatase

gene from tomato. The amplified and cloned 522 bp long fragment is not identical to

*LeMKP1*. Therefore, its corresponding gene is named *LeMKP2* (SEQ ID NO: 23 encoding SEQ ID NO: 24; 75% identity on the nucleotide level over the stretch of 196 bp of

*ZmMKP1* flanked by primers 1 and 3). The origins of all identified MAP kinase phosphatase homologous gene sequences are confirmed by Southern blot analysis.

The following Table shows an alignment of a continuous stretch of 312 amino acids of *AtMKP1* with the related amino acid sequence of *ZmMKP2*.

139	AtMKP1	KREKIAFFDKKCSKVAADHYVVGDAVAKDKSIKNNIGITHILNCGVFICP	188
24	ZmMKP2	RKDQIAFFDKKCSKVAADHVYLGDAVAKDKRDILFRKNGITHILNCGVFCP	73
189	AtMKP1	EYFKSDFCYRSIWLQDSPSEDITTSILYDVFDFEEDVREBQSGRIFFVHCQG	238
74	ZmMKP2	EYFKSDBLVYRTIWLQDSPFEDITTSILYDVFDFEEDVREBQSGRVLVHCQG	123
239	AtMKP1	VSRSTSLVLAIVLMMRBQGSFDDAFQYVKASRGIALDFMKGFAQCQLTCCQKR	288
124	ZmMKP2	VSRSTSLVLAIVLMMRBQGSFDDAFQYVKASRGIALDFMKGFAQCQLTCCQKR	173
289	AtMKP1	VHAFPLSPTSLRMYYKMSPHSPYDPLHLVPMQLNDPCFGSLDSRGAFIIQ	338
174	ZmMKP2	VHAIPLSPNSVLRMYYRMAPHSQYAPLHLVPMQLNDPSPATLDSRGAFIVH	223
339	AtMKP1	LPSTLYTWGRQCEETMEKDAKAAVCOIARYEKVEAPLWVVRBGEDEFVY	388
224	ZmMKP2	VLSSLYVWVGKCDPVMKEKDAKAAAFQVVRKYEKVGHIKVVREGLLEPQEF	273
389	AtMKP1	WDFAFASILLFMIGG.....SVTKVQFGDRKVDAYNLDFEIFFOKAIE	428
274	ZmMKP2	WDAFSSMPFNSDSNWTKISKDQIDSAKSDPGSRKNESYDADFELVYKAIT	323
429	AtMKP1	GGFVPTTLASSNNEHEHTLHPARE	450
324	ZmMKP2	GGVVPAPFSTSGAGDEHTLHPARE	345

## SEQUENCE LISTING

&lt;110&gt; Novartis AG

&lt;120&gt; Organic compounds

&lt;130&gt; Silencing Gene

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 26

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 6356

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 1

gttcaagggtg gttgtatctn cagtattaga aagagagctt ctagagagag tttctaattc 60  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 120  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 180  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 240  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 300  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 360  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 420  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 480  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 540  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 600  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 660  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 720  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 780  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 840  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 900  
aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag aagagagagag 960  
gattcatttaa aactacattaa ttatctcttc tccaataagt cgttaaaaac ttaacattaa caaaaagttc 1020  
aattcgcagcaa gaagagttta ttcaataagt ttcaataagt ttcaataagt ttcaataagt 1080  
aagagtgtrcaa acaacacctg ggatcacatat ttcatgacct aaaaatcgatat aacaacacct 1140  
aaggttaattca gtaattttga tggcttaaac aaatatatgt aaaaatatgt accattttga ccaatgagca 1200  
tttttttggaa ataaaactac tgcatacgac ctggttcaaac acatgaaactg ccaatgacctt 1260  
taaaaactacc ttttaaacct tttagacctt tttagacctt tttagacctt tttagacctt 1320  
aaaccataac attatgattc gtaatttcga gtgatgatgt tttagagtgta tgtgaatgat 1380  
aaccataacac acaatrtgtg atatatatc actcatacat atcaaatca aactagttta 1440  
tcgtttccaa acaatgattga taaaatgtaa ttactatca aaaccttaag ccagrtagtt 1500  
aattaaacaa aatataaatag tatatgtrtc atatttcttc ttcttctc ttcttctc 1560  
aattaatatt acaaacaaag caaaaaaaa gaataaataa aacaaactc attaactgtc 1620  
aatcaattat cagaggggaaa agttactaag aaatgtcaac caatgagtat catatcatc 1680  
atgccttttt agaccctctt tttaattca ttcaagattt gtcctgttta taaatttgaa 1740  
gttaaaattcc aaaaatgtaa attctacaat ttcttaacaa ttctctccca catatgtta 1800  
ttaaagattcc caattacttc gtttaatttc ttctctccca ttctctccca ttctctccca 1860  
ttaaagattcc caattacttc gtttaatttc ttcttaacaa ttctctccca catatgtta 1920  
atcatgatat attagatat attagatat attagatat attagatat attagatat 1980

**What is claimed is:**

1. A DNA comprising an open reading frame encoding a protein characterized by a component amino acid sequence having 40% or more identity with an aligned component sequence of SEQ ID NO: 3
2. The DNA according to claim 1 comprising an open reading frame encoding a plant MAP kinase phosphatase.
3. The DNA according to claim 1 wherein the open reading frame encodes an amino acid sequence selected from the group of amino acid sequences described in SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.
4. The DNA according to claim 1, wherein the open reading frame encodes a protein characterized by the amino acid sequence of SEQ ID NO: 3
5. The DNA according to claim 1 characterized by the nucleotide sequence of SEQ ID NO: 1
6. The DNA according to claim 1 wherein the open reading frame encodes a protein contributing to repair of DNA damage in a plant cell.
7. The DNA according to claim 1 wherein the open reading frame encodes a protein conferring hypersensitivity to treatment with methyl methanesulfonate (MMS).
8. The DNA according to claim 7 wherein the open reading frame encodes a protein conferring hypersensitivity to treatment with UV light or X-rays.
9. The DNA according to claim 7 wherein the open reading frame encodes a protein interfering with abscisic acid signal transduction.
10. The protein encoded by the open reading frame of any one of claims 1 to 9.
11. A method of producing DNA according to claim 1, comprising
  - screening a DNA library for clones which are capable of hybridizing to a fragment of the DNA defined by SEQ ID NO: 1, wherein said fragment has a length of at least 15 nucleotides;
  - sequencing hybridizing clones;
  - purifying vector DNA of clones comprising an open reading frame encoding a protein with more than 40% sequence identity to SEQ ID NO: 3
  - optionally further processing the purified DNA.
12. A polymerase chain reaction wherein at least one oligonucleotide used comprises a sequence of nucleotides which represents 15 or more basepairs of SEQ ID NO: 1.



gaataataaa tgrttagat agatgaat taataaaaa ataaagatga 2040  
gtataaact ttgtttct ttgttaaatg aaatatagaa aaatatcat 2100  
altacataa ttaataaa tgtcagat aaacgttagt atataaaac 2160  
agaaaaaaca gaattatca tcttgaca gaataaataa aaatgctga 2220  
ttcaaat ttcaagct tctataata agaatcaat ttctgtagc 2280  
ataataatca taataaat aatcgtatc atctataat actccaaa 2340  
cgaattatc ttaattatc ttgtttatc atgtttct ccaactcc 2400  
atcttct tcttct tcttct tcttct tcttct tcttct 2460  
ctctctc gaagaaaaa aaataaaga tcttctc taaatcttca 2520  
caatcgtaa gttcaaat tcttctc ccttctc tcttctc 2580  
ccaggaat ttctcaat ccttctc ccttctc ccttctc 2640  
ctgtttct ttcttag ttcttag ttcttag ttcttag ttcttag 2700  
ttcttttt ttcttag ttcttag ttcttag ttcttag ttcttag 2760  
gttatctc ttcttag ttcttag ttcttag ttcttag ttcttag 2820  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 2880  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 2940  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3000  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3060  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3120  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3180  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3240  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3300  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3360  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3420  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3480  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3540  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3600  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3660  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3720  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3780  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3840  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3900  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 3960  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4020  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4080  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4140  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4200  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4260  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4320  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4380  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4440  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4500  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4560  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4620  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4680  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4740  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4800  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4860  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4920  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 4980  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 5040  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 5100  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 5160  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 5220  
ttcttag ttcttag ttcttag ttcttag ttcttag ttcttag 5280

tacaatctgg tgaatccatc ttgtgtcagaa ttgtactcagc aaagatatcc cctatatagg 5340  
 taataatagc ctaaaccttg gaggcctcga taccctttag taattgttaa gtcctgcgtc 5400  
 tcaatcttgg tactgaactac ttaggacagt tatattcga agatgaaat tacaatctca 5460  
 gcaatctcat tactgaactac ttaggacagt tatattcga agatgaaat tacaatctca 5460  
 tatgtatttg ttattctct agtggttg atgttccaa cttgttaga tcccttctgt 5580  
 gtttgttaac gatagttg ttctctct ttgttaacaga ttgttaggaa atctgagat 5640  
 cagaaagagt ttctgaagac ttctgaagcgt ctataacac caccgcggaag cttacacat 5700  
 tcaatcttgg ttcttctac agatctctc gaagcgaac ttgttaacac ctttccatc ttcagacaga 5760  
 gtaatctc agcctcagtg aatatctc gaagcgaac ttgttaacac ctttccatc ttcagacaga 5820  
 gttgcatccg aatacaatcg ttgtcttg attatgttca atttgctatc gatcatgtat 5880  
 ttgttttaga ggaatgttgg acaaaaaa cttagagct cgaatgaacg gtttttaca 5940  
 aatctctca gacatattt gatgtgggg gaaacaaac tctgttaga ttgggaaaa acttctctc 6060  
 cttctctca tcaatgaaat atctcgaag gcaatttta gtatgacag attataaat 6120  
 ttgaagttta gggccttat gaggatttg tactaatttg tactaatttg tactaatttg 6180  
 tcaagacac aatagattt ttgttaacat tactaatttg tactaatttg tactaatttg 6240  
 atgttgaaat ctaattgaa attatgtta ttgtatttg tactaatttg tactaatttg 6300  
 atccaatctt gttttcgcg atagttctc ttcttttat ttcttttat ttcttttat 6356

<210> 2

<211> 3059

<212> DNA

<213> Arabidopsis thaliana

<400> 2

gtctctcc gcttcttc ccaatctc cttctctc tctctgaag aataataa 60  
 aaagatctaa cttgacgcg tctcttaac tcaatctc cagatgtgga tttaaacat 120  
 ctgattggag cttgattgat atgttagat aaagacccct gtttaccat gttatgttg 180  
 aacaaagtca ttgaagagc ttgaagcttt gatttgttaa ttgttaga ttgtatgttg 240  
 ctgatttttg gttgaatagg ttgaagcttt gatttgttaa ttgttaga ttgtatgttg 300  
 gttggaaagag aggtctcga ttgtgaatgat gaagcctcc cttgtctcaa ttgttaga 360  
 ttggtgtctg cttcttctg ttgtctcga ttgttaga ttgtatgttg ttgttaga 420  
 caaagcctga acattcttg ttgtctcga ttgttaga ttgtatgttg ttgttaga 480  
 cctttgacac cttgtctca tcaataacag ttgttaga ttgtatgttg ttgttaga 540  
 cctcttgcaa ttcttagag gacttagag ttgttaga ttgtatgttg ttgttaga 600  
 ggtgagttgg cttatccac aaacacccag ttgttaga ttgtatgttg ttgttaga 660  
 gattttctc caacgctgaag ttgttaga ttgtatgttg ttgttaga ttgttaga 720  
 aagattctc ttcttaga ttgttaga ttgtatgttg ttgttaga ttgttaga 780  
 ttgtcttggt ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 840  
 ttgtcttggt ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 900  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 960  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1020  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1080  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1140  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1200  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1260  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1320  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1380  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1440  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1500  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1560  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1620  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1680  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1740  
 ttgttaga ttgttaga ttgtatgttg ttgttaga ttgttaga ttgttaga 1800

actgctgc ctccttc ctcctccgat tctgtatgt caacaattc aggcataagc 1860  
 ttaagaagt tctcctcaatc ctcctgaagt tctgtcctga gacctctat tcccaagaa 1920  
 ctgacattgc ctaatttc cagcctatcc ctcctcctc cccaattc tcccaagaa 1980  
 tctcgtgtg tcaatattc tctcacaacg tcaacaataa gaagcctc acctctctc 2040  
 gctgagctga gagcagacct gaagagatct ctgaagtgc caggttgc tgattccaac 2100  
 agagcagcac ctgctttac tttacatccg gatgatagat atgacatagt ctccaatcg 2160  
 gaggtattga gaacggcgga tctatatcca ccaagtgtat gcaaggggac aagtgtagat 2220  
 tcaatttc aattatcc ttaatcagtt gcaagtaaac tgacagagac 2280  
 aatccggag gtatatacga tagctctggc cagccttag catgtcgtg gccaaagtatg 2340  
 gagatgatta caaaactgag cagagcttac ttagatcag aatcgttat agcaatccg 2400  
 tggccaagt atgctgtagg agaaacgggt agtagaat tgtacattg gatcggaaag 2460  
 tcatctctc tggtaaccaa ctgtctctca gtagatagca acaagcggc agaacactgtg 2520  
 gagaatgtg atgggtaca aatgggtgaa tccatttgt gtcagatggg ctggccaaa 2580  
 gataccctta taagatagt tagggaatc gagatcaga cagagtgtc agcaactgtg 2640  
 agcgcctat aacacccacc cgcaagctc acacattac tctgtttt ttccacagat 2700  
 tcttcaacc gcaacattc tccatttca gacagagtat tcatcagct caggaactc 2760  
 tgtccttag gaagttagt tgggggaa caactctc gtaagtggg gaaaaaactc 2820  
 ctctctc tttctac atgaaatcc tcaagggcca ttatagtat gatcagatata 2880  
 taaatgta agtttaggg cttatgagg attttagtg acttggtaca atgtttacat 2940  
 atacactag cagcacaata gattttgt aaactacat gttatcag gttaaaagtact 3000  
 atgtatgt tgaagtctaa ttgaagaat agttaatgat aaaaaaaaaa aaaaaaaa 3059

<210> 3  
 <211> 784  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 3  
 Met Val Gly Arg Glu Asp Ala Met Gly Asn Asp Glu Ala Pro Pro Gly 1  
 5  
 10  
 15  
 Ser Lys Lys Met Phe Trp Arg Ser Ala Ser Trp Ser Ala Ser Arg Thr 20  
 25  
 30  
 Ala Ser Gln Val Pro Gln Gly Asp Glu Gln Ser Leu Asn Ile Pro Cys 35  
 40  
 45  
 Ala Ile Ser Ser Gly Pro Ser Arg Arg Cys Pro Ala Ala Pro Leu Thr 50  
 55  
 60  
 Pro Arg Ser His His Asn Ser Lys Ala Arg Ala Cys Leu Pro Pro Leu 65  
 70  
 75  
 80  
 Gln Pro Leu Ala Ile Ser Arg Arg Ser Leu Asp Glu Trp Pro Lys Ala 85  
 90  
 95  
 Gly Ser Asp Asp Val Gly Glu Trp Pro His Pro Pro Thr Pro Ser Gly 100  
 105  
 110  
 Asn Lys Thr Gly Glu Arg Leu Lys Leu Asp Leu Ser Ser Thr Gln Gln 115  
 120  
 125  
 Arg Val Thr Asp Lys Ser Ser Gly Leu Ala Lys Arg Glu Lys Ile Ala 130  
 135  
 140

Phe Phe Asp Lys Glu Cys Ser Lys Val Ala Asp His Ile Tyr Val Gly 145  
 150 155  
 Gly Asp Ala Val Ala Lys Asp Lys Ser Ile Leu Lys Asn Asn Gly Ile 165  
 170 175  
 Thr His Ile Leu Asn Cys Val Gly Phe Ile Cys Pro Glu Tyr Phe Lys 180  
 185  
 Ser Asp Phe Cys Tyr Arg Ser Leu Trp Leu Glu Asp Ser Pro Ser Glu 190  
 195 200  
 Asp Ile Thr Ser Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val 205  
 210 215  
 Arg Glu Glu Ser Gly Arg Ile Phe Val His Cys Cys Glu Gly Val Ser 220  
 225 230 235  
 Arg Ser Thr Ser Leu Val Ile Ala Tyr Leu Met Trp Arg Glu Glu 240  
 245 250 255  
 Ser Phe Asp Asp Ala Phe Glu Tyr Val Lys Ser Ala Arg Gly Ile Ala 260  
 265 270  
 Asp Pro Asn Met Gly Phe Ala Cys Glu Leu Leu Cys Glu Lys Arg 275  
 280 285  
 Val His Ala Phe Pro Leu Ser Pro Thr Ser Leu Leu Arg Met Tyr Lys 290  
 295 300  
 Met Ser Pro His Ser Pro Tyr Asp Pro Leu His Leu Val Pro Lys Leu 305  
 310 315  
 Leu Asn Asp Pro Cys Pro Gly Ser Leu Asp Ser Arg Gly Ala Phe Ile 320  
 325 330 335  
 Ile Glu Leu Pro Ser Ala Ile Tyr Ile Trp Val Gly Arg Glu Cys Glu 340  
 345 350  
 Thr Ile Met Glu Lys Asp Ala Lys Ala Val Cys Glu Ile Ala Arg 355  
 360 365  
 Tyr Glu Lys Val Glu Ala Pro Ile Met Val Val Arg Glu Gly Asp Glu 370  
 375 380  
 Pro Val Tyr Tyr Trp Asp Ala Phe Ala Ser Ile Leu Pro Met Ile Gly 385  
 390 395 400  
 Gly Ser Val Ile Lys Val Glu Pro Gly Asp Arg Lys Val Asp Ala Tyr 405  
 410 415  
 Asn Leu Asp Phe Glu Ile Phe Glu Lys Ala Ile Glu Gly Gly Phe Val 420  
 425 430  
 Pro Thr Leu Ala Ser Ser Asn Asn Glu His Glu Thr His Leu Pro Ala

Arg Gln Asn Ser Trp Ser Ser Leu Lys Cys Lys Phe Ala Ser Arg Phe  
445  
440  
445  
455  
450  
460  
470  
475  
480  
485  
490  
495  
500  
505  
510  
520  
525  
535  
540  
550  
555  
560  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
630  
635  
640  
645  
650  
655  
660  
665  
670  
680  
685  
690  
695  
700  
710  
715  
720  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000



1 5 10

<210> 8  
<211> 4  
<212> PRT  
<213> Arabidopsis thaliana  
<400> 8  
Phe Val His Cys  
1

<210> 9  
<211> 6  
<212> PRT  
<213> Arabidopsis thaliana  
<400> 9  
Gln Gly Val Ser Arg Ser  
1  
5

<210> 10  
<211> 5  
<212> PRT  
<213> Arabidopsis thaliana  
<400> 10  
Tyr Phe Lys Ser Asp  
1  
5

<210> 11  
<211> 23  
<212> DNA  
<213> Artificial Sequence  
<220>  
<223> Description of Artificial Sequence: degenerate  
primer

<400> 11  
aayaaaygma thacacayac hyt

<210> 12  
<211> 24  
<212> DNA  
<213> Artificial Sequence  
<220>  
<223> Description of Artificial Sequence: degenerate  
primer  
<400> 12





65

<210> 17  
<211> 196  
<212> DNA  
<213> Lycopersicon esculentum

<400> 17  
aaactgtta ggttagtt gtcctgaata ctttaaggat gaccttgat acaagacact 60  
ttgctgcag gataggccca ctgaggacat cacaagatat cttatgatg tcttgatta 120  
cttgaagat gttcatgaac aagtgaggag tgccttgta cactgcttcc aggggtgtc 180  
ccgatacagc tctctg 196

<210> 18  
<211> 65  
<212> PRT  
<213> Lycopersicon esculentum

<400> 18  
Asn Cys Val Gly Phe Ser Cys Pro Gln Tyr Phe Lys Asp Asp Leu Val 1  
5  
Tyr Lys Thr Leu Trp Leu Gln Asp Ser Pro Thr Gln Asp Ile Thr Ser 20  
30  
Ile Leu Tyr Asp Val Phe Asp Tyr Phe Gln Asp Val His Gln Gly 35  
40  
Gly Ser Val Phe Val His Cys Phe Gln Gly Val Ser Arg Ser Ala Ser 50  
60  
Leu 65

<210> 19  
<211> 196  
<212> DNA  
<213> Zea mays

<400> 19  
caactgctg gcttcgtct gcccgaaata ctcaagtc gaccttct accgacact 60  
ctgctcag gacagcccca ccgaggacac caccagcatc cttaacgacg tgttgatta 120  
cttgagagc gtcaaggagc aggtggccg cgtgttctg catgctgcc aggggtgtc 180  
gctccacg cctctg 196

<210> 20  
<211> 65  
<212> PRT  
<213> Zea mays

<400> 20  
Asn Cys Met Gly Phe Val Cys Pro Lys Tyr Phe Lys Ser Asp Leu Val 1  
5  
10  
15

Tyr Arg Thr Leu Trp Leu Asp Ser Pro Thr Glu Asp Thr Ser 20  
 Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val Arg Glu Glu Gly 35  
 Gly Arg Val Phe Val His Cys Cys Glu Gly Val Ser Arg Ser Thr Pro 50  
 40 45 55 60

Leu 65

<210> 21  
 <211> 2450  
 <212> DNA  
 <213> Zea mays

<400> 21

ggcggtgcct ccccgcccaa gccggggag gggctccgac tcgaactctc ctcgctccgg 60  
 tcgcaggggc gcaaggacca gatcgccctc ttcgacaagg agtgcctcaa agtgcctccg 120  
 cagrtctaac tcggcgcgga cgcgtccgac aagaaacggg acatccctcag gaaagaaagg 180  
 atcaccccaa tgcctcaactg tgtggggcctt gtcctggccgg agtactccta gtcggacct 240  
 gtctaccga cccctctggc gcaggagcag cccaaccggg acatccctcag gaaagaaagg 180  
 tgcagggggc tgcagggggc cagrtctcag gacgtctcag gacgagggcg gccgctgct 360  
 gacgtgttg actactcga gacgtctcag gacgtctcag gacgtctcag gacgtctcag 420  
 cagagctcg atgacgctc cagrtctcag aaggtctgac gggggatcgc aaatcccaac 480  
 atggccttg catgcccagc tcctcagtgcc caggaagcttg tgcattgctg tcctctgtca 540  
 ccaatrcag tgcctcaggg gtcctcaggg gtcctcaggg gtcctcaggg gtcctcaggg 600  
 ttgtgtccca aaatgctcaa tgaacctcc ccaagcccaac ttgactcag aggcgcgttc 660  
 atgtctcgtc gctcctatgc tgggttgga tgaagtgtga tccggtaatg 720  
 gaaaggatg caaagctgc tgcgttcag gtagtgaggt atgaagaagt gcaaggggcac 780  
 atcaagtg tgaagaagg tcaggagcg caggagttct gggatgcat tccaagt 840  
 ccaactaat cagatagtaa tacaagatc agcaaggacc agatgcatc agcatccaag 900  
 agtgaaccag gaagccggaa aaatgagtc tatgagctg atttgagc tgtctcaaa 960  
 gcaatrcactg ggggagtggt cccgtgcatt tcgaactctg gggctgtg tgaagcccat 1020  
 cttccagcta gagaagtag ctggagttca atggatttt atccacgggt acaaacacct 1080  
 gctcgtgttt atcagatcc tgcctcaatg aaggtatttg atccacgggt acaaacacct 1140  
 gctgctgagg catcaacctc acctccttc cttcctcaa gctcctatc atcggtatca 1200  
 agtgcagct cgaagtatag ttcagactca cccctctat caactacaac tggctcctca 1260  
 ccatcattg gctcctcgcc tgcctcatc aatctgacac atgctcttgg gccatcatcc 1320  
 agtctctccc ttctcatc atctaataa ggaagcttcaa agcctcttg catggaatca 1380  
 atacaacttc cttccaaagc gcatctcata gcagaaaggag gaggaggtct cacactctca 1440  
 aagctaacat cttcccaaaa ggaatcttga ttgccacca ggtgtgcgtc tatgttatg 1500  
 ccaaccaagg agctcatcag tatctcgaggg aaccgaggaatg cctcagataa tagtacaat 1560  
 tgggttaaac agctgacatg tgaatttgc actgacagta actcagaaat ctgcagtaat 1680  
 gcaacaactg taatcagatc atggtccagc taactaatg atggtccagc tgcacggcaa 1740  
 gatcttgacc cgaagtcgt tttaatttt gtaactcga atggtccagc gagggagaa 1800  
 gcaagttaaaa tgggtatgt atgggtatgg aggggtatgg aggtgtatg 1860  
 tccgtctcag aggttttcaa aggttttcaa aggttttcaa aggttttcaa aggttttcaa 1920  
 cttccctcca aggttttcaa aggttttcaa aggttttcaa aggttttcaa aggttttcaa 1980  
 agttaaactt aggcagtagc tatcagatata atgttagtg ctaaaccaac tcaacgaagg 2040

catgccctcc agcatcagtc ggtaccgagtg atgtcagcgg aggtataaag ccacagccat 2100  
 tcccttgaaac ataataagct accaaacagat tccgttctgc aactgcgcct catgatctat 2160  
 attttgtcca gatggcagga ggcgtgccatg ggcgttgtat cggttgcgaa ttagcactcg 2220  
 tgggttagga gcaatccggcc gatccggtgt atatatccg ctcctccctga atgtaagctc 2280  
 agatactggg agctgggtg tcgacagtta cttttagcc taacatctc tgtacatctc 2340  
 tgaagggaac agagtgtaa tcctttgac tatgtaaatg gctccatgg tcataacttc 2400  
 taaccatgt ggaactcagt tgcctagctgg taaaaaaaaa aaaaaaaaaa 2450

<210> 22  
 <211> 661  
 <212> PRT  
 <213> Zaa may

<400> 22  
 Gly Gly Ser Ser Pro Ala Lys Pro Gly Glu Gly Leu Arg Leu Asp Leu 1  
 5 10 15

Ser Ser Leu Arg Ser Gln Gly Arg Lys Asp Gln Ile Ala Phe Asp 20  
 25 30

Lys Glu Cys Ser Lys Val Ala Asp His Val Tyr Leu Gly Gly Asp Ala 35  
 40 45

Val Ala Lys Asn Arg Asp Ile Leu Arg Lys Asn Gly Ile Thr His Val 50  
 55 60

Leu Asn Cys Val Gly Phe Val Cys Pro Glu Tyr Phe Lys Ser Asp Leu 65  
 70 75 80

Val Tyr Arg Thr Leu Trp Leu Gln Asp Ser Pro Thr Glu Asp Ile Thr 85  
 90 95

Ser Ile Leu Tyr Asp Val Phe Asp Tyr Phe Glu Asp Val Arg Glu Gln 100  
 105 110

Gly Gly Arg Val Leu Val His Cys Cys Gln Gly Val Ser Arg Ser Thr 115  
 120 125

Ser Leu Val Ile Ala Tyr Leu Met Trp Arg Glu Gly Gln Ser Phe Asp 130  
 135 140

Asp Ala Phe Gln Phe Val Lys Ala Ala Arg Gly Ile Ala Asn Pro Asn 145  
 150 155 160

Met Gly Phe Ala Cys Gln Leu Leu Gln Cys Gln Lys Arg Val His Ala 165  
 170 175

Ile Pro Leu Ser Pro Asn Ser Val Leu Arg Met Tyr Arg Met Ala Pro 180  
 185 190

His Ser Gln Tyr Ala Pro Leu His Leu Val Pro Lys Met Leu Asn Asp 195  
 200 205

Pro Ser Pro Ala Thr Leu Asp Ser Arg Gly Ala Phe Ile Val His Val 210  
 215 220

Leu Ser Ser Leu Tyr Val Trp Val Gly Met Lys Cys Asp Pro Val Met 225  
 230 235 240  
 Glu Lys Asp Ala Lys Ala Phe Gln Val Val Arg Tyr Glu Lys 245  
 250 255  
 Val Gln Gly His Ile Lys Val Val Arg Glu Gly Leu Glu Pro Gln Glu 260  
 265 270  
 Phe Trp Asp Ala Phe Ser Ser Met Pro Pro Asn Ser Asp Ser Asn Thr 275  
 280 285  
 Lys Ile Ser Lys Asp Gln Ile Asp Ser Ala Ser Lys Ser Asp Pro Gly 290  
 295 300  
 Ser Arg Lys Asn Glu Ser Tyr Asp Ala Asp Phe Glu Leu Val Tyr Lys 305  
 310 315  
 Ala Ile Thr Gly Gly Val Val Pro Ala Phe Ser Thr Ser Gly Ala Gly 325  
 330 335  
 Asp Glu Thr His Leu Pro Ala Arg Glu Ser Ser Trp Ser Leu Leu Arg 340  
 345 350  
 His Lys Phe Ile Ser Arg Ser Leu Ala Arg Val Tyr Ser Asp Ser Ala 355  
 360 365  
 Leu Met Lys Asp Phe Asp Pro Arg Val Gln His Leu Ala Ala Glu Ala 370  
 375 380  
 Ser Thr Ser Pro Pro Phe Leu Ser Ser Ser Leu Ser Ser Ser Asp Ser 385  
 390 395 400  
 Ser Val Ser Ser Lys Tyr Ser Ser Asp Ser Pro Ser Leu Ser Pro Thr 405  
 410 415  
 Thr Gly Ser Pro Pro Ser Phe Gly Leu Ser Pro Ala Ser Ser Asn Leu 420  
 425 430  
 Thr His Ala Leu Val Pro Ser Ser Arg Ser Pro Leu Ser Gln Ser Ser 435  
 440 445  
 Asn Glu Gly Ala Ser Lys Pro Ser Gly Met Glu Ser Ile His Ser Pro 450  
 455 460  
 Ser Lys Thr Ser Ser Ile Ala Glu Arg Arg Gly Gly Phe Thr Leu Leu 465  
 470 475 480  
 Lys Leu Pro Ser Leu Gln Lys Asp Leu Val Leu Pro Pro Arg Val Pro 485  
 490 495  
 Ser Ile Val Leu Pro Pro Arg Ala Pro Ser Ser Ile Arg Arg Thr Glu 500  
 505 510

Asn Cys Val Gly Phe Val Cys Pro Glu Tyr Phe Lys Ser Asp Phe Val  
1 5 10 15

<400> 24  
<212> PRT  
<211> 174  
<210> 24

aactgtgtgtg ggtttgtatg ccagagtat ttcaagtcg atttcgtata ccggaacttg 60  
tggttgcagg atagcccatc agaagatat actagatat tctatgatgt ttctgaactac 120  
tttgaagatg tcaaggagca acatgggaag gttttgttc atgtctgcca aggggtctct 180  
cggtcaacct cgttggttat tgcctatcgt atgtggagag aaggacaag ttcttgatgat 240  
gaccttgagt atgtaaaagc agcaaggagt atgtcgagtc caaatatggg ttctgcttct 300  
catttatcac aatgccaaaa aagggttcat gcttctctt tgaagcccaag ttcatlata 360  
aggatgtaca gacttgccac tcaatccaac tagatctctt tgcattctct cccaaaaatg 420  
ttaaatgatc cctcacccggc agcatatgat tctagaggtg cattatat acacataact 480  
tcatcggtat atgtatgat tggtaagaaa tgtgaagcaa tc 522

<400> 23  
<213> Lycopersicon esculentum  
<212> DNA  
<211> 522  
<210> 23

Leu Leu Asn Val Ser 660  
Leu Pro Val Lys Val Phe Lys Glu His Glu Ala Glu Asn Leu Glu 645  
Gln Gln Val Thr Ser Asp Phe Leu His Leu Lys Gly Leu Ser Asn Val 625  
Val Gly Gly Glu Asn Glu Ser Ser Lys Ser Val Asp Ser Val Asp Trp 610  
Ser Asn Ala Ile Arg Arg Gly Glu Ala Val Lys Met Val Tyr Val Trp 595  
Phe Ala Arg Lys Asp Leu Asp Pro Lys Ser Val Leu Ile Phe Val Thr 580  
Ala Gln Leu Val Val Tyr Gln Trp Pro Ser Met Gly Lys Leu Thr 565  
Thr Arg Leu Ile Glu Arg Thr Asp Ser Asn Ser Glu Val Cys Ser Asn 545  
Phe Cys Ser Glu Lys Cys Thr Gly Asn Ser Leu Ser Ser His Ser Glu 530  
Asp Ala Ser Asp Asn Ser Thr Asn Gly Val Lys Gln Leu Thr Ser Glu 515  
512 525

Tyr Arg Thr Leu Trp Leu Gln Asp Ser Pro Ser Gln Asp Ile Thr Ser  
 20 25 30  
 Ile Leu Tyr Asp Val Phe Asp Tyr Phe Gln Asp Val Arg Gln His  
 35 40 45  
 Gly Lys Val Phe Val His Cys Gln Gly Val Ser Arg Ser Thr Ser  
 50 55 60  
 Leu Val Ile Ala Tyr Arg Met Trp Arg Gln Gly Gln Ser Phe Asp Asp  
 65 70 75 80  
 Ala Phe Gln Tyr Val Lys Ala Ala Arg Gly Ile Ala Asp Pro Asn Met  
 85 90 95  
 Gly Phe Ala Cys Gln Leu Leu Gln Cys Gln Lys Arg Val His Ala Ser  
 100 105 110  
 Pro Leu Ser Pro Ser Ser Leu Leu Arg Met Tyr Arg Val Ala Pro His  
 115 120 125  
 Ser Pro Tyr Asp Pro Leu His Leu Val Pro Lys Met Leu Asn Asp Pro  
 130 135 140  
 Ser Pro Ala Ala Leu Asp Ser Arg Gly Ala Phe Ile His Ile Pro  
 145 150 155 160  
 Ser Ser Val Tyr Val Trp Ile Gly Lys Lys Cys Gln Ala Ile  
 165 170

<210> 25  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: degenerate  
 primer  
 <400> 25  
 gcngcycttc c  
 21  
 <210> 26  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: degenerate  
 primer  
 <400> 26  
 ytcnchngcn ggnarrtngn tyc  
 24

# INTERNATIONAL SEARCH REPORT

Internal Application No

PCT/EP 99/05413

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N9/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

X	WO 94 23039 A (CANCER RES INST ROYAL MARSHALL CHRISTOPHER JOHN (GB); ASHWORTH AL) 13 October 1994 (1994-10-13) figure 7D & DATABASE GENE, online!	1-12
X	Accession no. R63602, 9 May 1995 (1995-05-09) abstract	1-12
X	DATABASE SWISSPROT, online! Accession no. P28562, 1 December 1992 (1992-12-01) KEYES S.M. AND EMSLIE E.A.: "DUAL SPECIFICITY PROTEIN PHOSPHATASE 1" XP002131991 abstract	1-12
	---	
	---	
	---	

<input checked="" type="checkbox"/>	Further documents are listed in the continuation of box C.
<input checked="" type="checkbox"/>	Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but
"E" earlier document but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the
"O" document referring to an oral disclosure, use, exhibition or other means	"Z" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the
"P" document published prior to the international filing date but later than the priority date claimed	"a" document member of the same patent family in the art.

Date of the actual completion of the international search	Date of mailing of the international search report
1 March 2000	14/03/2000

Name and mailing address of the ISA European Patent Office, P.B. 5018 Patentkanal 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016	Authorized officer Chakravarty, A
--	--------------------------------------

Form PCT/ISA/210 (second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 99/05413

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE SWISSPROT 'online' Accession no. P28563, 1 December 1992 (1992-12-01) CHARLES C.H. ET AL.: "DUAL SPECIFICITY PROTEIN PHOSPHATASE 1 (EC 3.1.3.48) (EC 3.1.3.16)" XP002131992 cited in the application abstract</p>	1-12
P, X	<p>DATABASE EMBL 'online' EBI accession no. AL085836, 28 June 1999 (1999-06-28) SALANOUBAT M. ET AL.: "17 end of BAC F1115 of IGF library from strain Columbia of Arabidopsis thaliana" XP002131993 abstract</p>	1-12



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/05413

Information on patent family members

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
---	---------------------	----------------------------	---------------------

WO 9423039	A	13-10-1994	AU 677834 B	08-05-1997
			AU 6382394 A	24-10-1994
			CA 2157774 A	13-10-1994
			EP 0703984 A	03-04-1996
			JP 9501302 T	10-02-1997
			US 5958721 A	28-09-1999
			AU 696939 B	24-09-1998
			AU 1586195 A	29-08-1995
			CA 2182967 A	17-08-1995
			EP 0742827 A	20-11-1996
			WO 9521923 A	17-08-1995
			JP 9508795 T	09-09-1997

